References – kesemek

Akagi, T., Katayama-Ikegami, A., & Yonemori, K. (2011). Proanthocyanidin biosynthesis of persimmon (Diospyros kaki Thunb.) fruit. Scientia Horticulturae, 130(2), 373 380.

Chang, Y.-L., Lin, J.-T., Lin, H.-L., Liao, P.-L., Wu, P.-J., & Yang, D.-J. (2019). Phenolic compositions and antioxidant properties of leaves of eight persimmon varieties harvested in different periods. Food Chemistry, 289, 74-83.

Gutiérrez-del-Río, I., Fernández, J., & Lombó, F. (2018). Plant nutraceuticals as antimicrobial agents in food preservation: Terpenoids, polyphenols and thiols. International journal of antimicrobial agents, 52(3), 309-315.

Heo, B.-G., Park, Y.-J., Kim, T.-C., Kim, H.-J., Park, S.-M., Jang, H.-G., Kim, K.-S., Lee, K.-D., & Yun, J.-G. (2009). Chemical characteristics, antimicrobial activity and dyeability of Gyeongsanbansi (Persimmon kaki) unripe juice extraction and fermented liquor. Korean Journal of Plant Resources, 22(5), 438-445.

Kamimoto, M., Nakai, Y., Tsuji, T., Shimamoto, T., & Shimamoto, T. (2014). Antiviral effects of persimmon extract on human norovirus and its surrogate, bacteriophage MS2. Journal of Food Science, 79(5), M941-M946.

Ketnawa, S., Hamanaka, D., Wongpoomchai, R., Ogawa, Y., & Setha, S. (2021). Low intensity of high pressure processing increases extractable recovery of polyphenols and antioxidant activities of non-astringent persimmon fruit. Lwt, 151, 112162.

Kim, H.-W., Lee, C.-H., & Lee, H.-S. (2009). Antibacterial activities of persimmon roots derived materials and 1, 4-naphthoquinone's derivatives against intestinal bacteria. Food Science and Biotechnology, 18(3), 755-760.

Liu, M., Yang, K., Wang, J., Zhang, J., Qi, Y., Wei, X., & Fan, M. (2019). Young astringent persimmon tannin inhibits methicillin-resistant Staphylococcus aureus isolated from pork. Lwt, 100, 48-55.

Moreno-Chamba, B., Salazar-Bermeo, J., Martínez-Madrid, M. C., Lizama, V., Martín Bermudo, F., Berná, G., Neacsu, M., Saura, D., Martí, N., & Valero, M. (2022). Bound galloylated compounds in persimmon upcycled dietary fiber modulate microbial strains associated to human health after in vitro digestion. Lwt, 156, 113011.

Pisoschi, A. M., Pop, A., Georgescu, C., Turcuș, V., Olah, N. K., & Mathe, E. (2018). An overview of natural antimicrobials role in food. European journal of medicinal chemistry, 143, 922-935.

Redpath, S., & George, A. (2008). Health and medicinal benefits of persimmon fruit: A review. Health and Medicinal Benefits of Persimmon Fruit, 1000-1006.

Tessmer, M. A., Besada, C., Hernando, I., Appezzato-da-Gloria, B., Quiles, A., & Salvador, A. (2016). Microstructural changes while persimmon fruits mature and ripen. Comparison between astringent and non-astringent cultivars. Postharvest Biology and Technology, 120, 52-60.

Villalobos-Delgado, L. H., Nevárez-Moorillon, G., Caro, I., Quinto, E. J., & Mateo, J. (2019). Natural antimicrobial agents to improve foods shelf life. In Food quality and shelf life (pp. 125-157). Elsevier.